



Unhealthy Alcohol and Drug Use

Adult, Primary Care, Clinical Practice Guideline Pocket Guide

UNHEALTHY ALCOHOL AND DRUG USE GUIDELINE USE

Full Dartmouth-Hitchcock Unhealthy Alcohol and Drug Use Adult, Primary Care Clinical Practice Guideline

http://sitefinity.hitchcock.org/intranet/docs/default-source/d-h-knowledge-map-documents/uadu-guideline-final_2017.pdf?sfvrsn=8

Unhealthy Alcohol and Drug Use Adult, Primary Care Clinical Practice Guideline Brief

http://sitefinity.hitchcock.org/intranet/docs/default-source/d-h-knowledge-map-documents/sud_brief_2_6_16.pdf?sfvrsn=2

Guideline Adoption Statement Source Documents:

[MA-SBIRT \(Massachusetts Screening, Brief Intervention and Referral to Treatment\) Clinician's Toolkit²](#)

[Helping Patients Who Drink Too Much: A Clinician's Guide¹](#)

[Facing Addiction in America: The Surgeon General's Report on Alcohol, Drugs and Health¹](#)

D-H Depression Management Guideline

http://sitefinity.hitchcock.org/intranet/docs/default-source/d-h-knowledge-map-documents/depression_cpg_final.pdf?sfvrsn=2

D-H Behavioral Health Integration into Primary Care Model Guideline

<http://sitefinity.hitchcock.org/intranet/docs/default-source/d-h-knowledge-map-documents/behavioral-health-integration-guideline-final.pdf?sfvrsn=2>

Definitions:

Risky Use: consumption of amounts that increase the likelihood of health consequences.


Risky Drinking¹:

For healthy men up to age 65:

- more than 4 drinks in a day or
- more than 14 drinks in a week

For all healthy women and healthy men over age 65:

- more than 3 drinks in a day or
- more than 7 drinks in a week



Copyright, citation, use, and adoption limitation/instructions:

May not be reproduced, distributed or modified for sale. May not be loaded into software platform outside of D-H in whole or in part without explicit permission.

If modified for local use, cite as:

Adapted from Dartmouth-Hitchcock Knowledge Map™ Unhealthy Alcohol and Drug Use, Adult, Primary Care Clinical Practice Guideline. Copyright 2017.

Pathways & Guidelines: Clinical Practice Guideline and pathways are designed to assist clinicians by providing a framework for the evaluation and treatment of patients. This Clinical Practice Guideline outlines the preferred approach for most patients. It is not intended to replace a clinician's judgment or to establish a protocol for all patients. It is understood that some patients will not fit the clinical condition contemplated by a guideline and that a guideline will rarely establish the only appropriate approach to a problem.

Unhealthy Alcohol and Drug Use, Adult, Primary Care Clinical Practice Guideline Pocket Guide

Copyright © 2017 Dartmouth-Hitchcock Knowledge Map™. Updated: January 2017

Algorithm: Recognizing and Managing Alcohol and Substance Use Disorders in Primary Care

